

REPORT DOCUMENTATION PAGE			Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>				
1. REPORT DATE (DD-MM-YYYY) February 2013		2. REPORT TYPE Viewgraph		3. DATES COVERED (From - To) February 2013- April 2013
4. TITLE AND SUBTITLE Thermal Decomposition Mechanisms of Ionic Liquids by Direct Dynamics Simulations and Vacuum Ultraviolet Photoionization Mass Spectrometry			5a. CONTRACT NUMBER FA9300-06-C-0023	
			5b. GRANT NUMBER	
			5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) S. D. Chambreau, J. Liu, G. L. Vaghjiani, C. J. Koh, A. Golan, O. Kostko, S. R. Leone			5d. PROJECT NUMBER	
			5e. TASK NUMBER	
			5f. WORK UNIT NUMBER 23030423	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQRP 10 E. Saturn Blvd. Edwards AFB CA 93524-7680			8. PERFORMING ORGANIZATION REPORT NO.	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) Air Force Research Laboratory (AFMC) AFRL/RQR 5 Pollux Drive Edwards AFB CA 93524-7048			10. SPONSOR/MONITOR'S ACRONYM(S)	
			11. SPONSOR/MONITOR'S REPORT NUMBER(S) AFRL-RQ-ED-VG-2013-042	
12. DISTRIBUTION / AVAILABILITY STATEMENT Distribution A: Approved for Public Release; Distribution Unlimited. PA#13183				
13. SUPPLEMENTARY NOTES Viewgraph for the American Chemical Society Meeting, New Orleans, LA, 6-11 April 2013.				
14. ABSTRACT Due to the unusually high heats of vaporization of room-temperature ionic liquids (RTILs), volatilization of RTILs through thermal decomposition and vaporization of the decomposition products can be significant. In complex molecules like RTILs, use of chemical intuition to predict reaction pathways can prove unreliable, especially when the internal energy content is high. There may be concerted reactions that are difficult to predict, and once energy stored in the molecule begins to release, the system does not necessarily follow the minimum energy reaction path, i.e., the subsequent behavior is controlled by dynamics. A useful approach to treating such a system is quasi-classical, direct dynamics trajectory simulations, where the motion of the molecule is followed, allowing the molecule to "show us" what the preferred reaction pathways are. The direct dynamics method dispenses with the potential energy surface. Instead, it calculates the energies, force constants, and Hessian "on the fly" using quantum chemistry methods. This method becomes computationally attractive when the dimensionality of the system increases, particularly for RTILs, which typically contain 10 or more heavy atoms. Dynamics simulations allows the partitioning of the energy generated by exothermic reactions into vibrational, rotational, and translational degrees of freedom, thereby increasing the chance of locating new reaction pathways in the system. In addition, by following the variation of the potential energy during the trajectory rather than relying on intuition, we can identify better geometries for TS searching. Results obtained from these direct dynamics simulations are compared to and consistent with the gaseous products detected experimentally via tunable vacuum ultraviolet photoionization mass spectrometry performed at the Chemical Dynamics Beamline 9.0.2 at the Advanced Light Source. The likely reaction mechanisms in the thermal decomposition of RTILs are discussed in this work.				
15. SUBJECT TERMS				
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT  SAR	18. NUMBER OF PAGES  44
a. REPORT Unclassified	b. ABSTRACT Unclassified	c. THIS PAGE Unclassified		19a. NAME OF RESPONSIBLE PERSON Tommy Hawkins
				19b. TELEPHONE NO (include area code) 661-525-5449



# Thermal decomposition mechanisms of ionic liquids by direct dynamics simulations and vacuum ultraviolet photoionization mass spectrometry

**Steve Chambreau**

*American Chemical Society Spring National Meeting  
New Orleans, Louisiana  
April 10, 2013*

*ERC, Inc.  
Aerospace Systems Directorate  
Propellants Branch  
Air Force Research Laboratory  
AFRL/RQRP  
10 E Saturn Blvd  
Edwards AFB, CA 93524*

**Tel:** 661 275 6176

**Fax:** 661 275 5471

**Email:** [steven.chambreau.ctr@edwards.af.mil](mailto:steven.chambreau.ctr@edwards.af.mil)






# Outline



- Direct dynamics and RRKM modeling of:
  - 1,5-dinitrobiuret (DNB) pyrolysis.
  - EMIM<sup>+</sup>dca<sup>-</sup>, EMMIM<sup>+</sup>dca<sup>-</sup>
- IL vapor studies:
  - Photoionization of vaporized ILs.
- IL aerosols + HNO<sub>3</sub>:
  - Diffusion-limited reactivity: MM
- QM/MM reactive scattering:
  - hyperthermal O(<sup>3</sup>P) off IL surface



# What is *hypergolicity*??!



*J. Phys. Chem. A*, **2008**, 112 (34), pp 7816–7824  
DOI: 10.1021/jp8038175





# Motivation

- Replacement for monomethylhydrazine (MMH) +  $\text{N}_2\text{O}_4$  (highly volatile and toxic!!)






# Ionic Liquids

- Molten salts with m.p.  $\leq 100$  °C.
- Asymmetric ions with diffuse charge distributions.
- $C^+A^-$  :  $10^{18}$  possible combinations of cations and anions.



# Background




- Dicyanamide-based RTILs first hypergolic ILs discovered with  $\text{HNO}_3$ .






# Gas Phase Decomposition of DNB





Geith, J., Holl, G., Klapötke, T. M., Weigand, J. J., *Combust. Flame*, **2004**, 139, 358-366

Frame #: 1



Frame #: 1





# RRKM Results

Temp / K	750	1000	1250	1500	1750	2000
density of states ( $1/\text{cm}^{-1}$ )						
DNB	$1.95 \times 10^{18}$	$2.34 \times 10^{22}$	$5.98 \times 10^{25}$	$4.94 \times 10^{28}$	$1.71 \times 10^{31}$	$2.92 \times 10^{33}$
twisted DNB	$1.90 \times 10^{18}$	$2.33 \times 10^{22}$	$6.04 \times 10^{25}$	$5.04 \times 10^{28}$	$1.75 \times 10^{31}$	$3.02 \times 10^{33}$
unimolecular rates ( $\text{s}^{-1}$ ) <sup>a</sup>						
$k_1$	0 (0)	0.6 (3.6)	$1.8 \times 10^4$ ( $1.8 \times 10^5$ )	$2.1 \times 10^6$ ( $2.7 \times 10^7$ )	$3.9 \times 10^7$ ( $5.8 \times 10^8$ )	$2.9 \times 10^8$ ( $4.7 \times 10^9$ )
$k_2$	8.5	$9.7 \times 10^5$	$7.9 \times 10^7$	$1.0 \times 10^9$	$5.5 \times 10^9$	$1.8 \times 10^{10}$
$k_{31}$	$3.4 \times 10^9$	$1.5 \times 10^{10}$	$3.5 \times 10^{10}$	$6.0 \times 10^{10}$	$8.8 \times 10^{10}$	$1.2 \times 10^{11}$
$k_{31}$	$2.4 \times 10^{11}$	$3.2 \times 10^{11}$	$3.8 \times 10^{11}$	$4.4 \times 10^{11}$	$4.8 \times 10^{11}$	$5.1 \times 10^{11}$
$k_{32}$	0	$1.5 \times 10^{-2}$	$2.1 \times 10^4$	$7.9 \times 10^6$	$2.9 \times 10^8$	$3.4 \times 10^9$
branching ratios (%) <sup>a</sup>						
path 1	0	0	0 (0.3)	0.2 (2.6)	0.7 (9.1)	1.3 (18.0)
path 2	100	100	100 (99.7)	99.0 (96.7)	94.3 (86.3)	83.3 (69.2)
path 3	0	0	0	0.8 (0.8)	5.0 (4.5)	15.4 (12.8)

() = loose transition state for path 1



# IL Vaporization

- Strasser, Armstrong, indicate ion pairs by detection of intact cation:<sup>1</sup>
  - **Ionic Liquid +  $\Delta \rightarrow \text{C}^+\text{A}^-_{(\text{g})}$**
  - **$\text{C}^+\text{A}^-_{(\text{g})} + h\nu \rightarrow \text{C}^+\text{A} + \text{e}^- \rightarrow \text{C}^+ + \text{A} + \text{e}^-$**
- Kelkar and Maginn: ion pairs more energetically favorable than clusters of ion pairs and non-neutral clusters.<sup>2</sup>


<sup>1</sup> *J. Phys. Chem. A*, 2007, **111**, 3191-3195; *PCCP*, 2007, **9**, 982-990.

<sup>2</sup> *J. Phys. Chem. B*, 2007, **111**, 9424-9427.





# ALS: Chemical Dynamics Beamline





7.4-15.0 eV photons, 0.025 eV resolution




# Experimental

IL effusive source:







Frame #: 1



Frame #: 1



Frame #: 1








# EMIM<sup>+</sup>dca<sup>-</sup> MD results

MD reaction	# of trajectories	branching ratio (%)
C2 H+ xfer to dca (term)	16	25
C2H5 dissoc	11	17
NCN + CN	8	13
dca dissoc	4	6
NR	4	6
dca abstract H from CH3 --> Hdca (term)	3	5
C2 H+ <-> dca (central)	2	3
CH3 dissoc	2	3
Et: CH3CH2- + dca --> CH2CH2 + Hdca (term)	2	3
H2 elim from -CH3	2	3
NCN-CN elongation	2	3
-C2H5 + dca <--> -C2H4 + Hdca	1	2
C2H5 abstraction by dca (term)	1	2
C2H5 and dca dissoc	1	2
C4H xfer to N3 and elim HC4C5	1	2
CH3 abstraction by dca (central)	1	2
CH3 abstraction by dca (term)	1	2
Et: H2 + dca --> HCN + HNCN	1	2
total=	63	




# PCM Model for ILs

- Truhlar: SMD-GIL (generic ionic liquid)\*
  - benchmark to experimental  $\Delta G_{\text{solv}}$  for ILs.



\*J. Phys. Chem. B **2012**, 116, 9122–9129







+ HCN



$$\begin{aligned} &\text{Mass 150} \\ &= (111 - 1) + 40 \end{aligned}$$




# addition-elimination mechanism















# photoionization efficiency of ion pair



J. Phys. Chem. Lett., **3**, 2910 (2012)

IP (M06/6-31+G(d,p))  
 $7.3 \pm 0.2$  eV



# Aerosol reactivity




- RTILs very low vapor:
  - Aerosols are liquid droplets suspended in gas phase.



# Monitoring isolated ion pairs

- Use it **to study reaction kinetics** of ‘hypergolic ionic liquid reaction with an oxidizer’


Hypergolic  
Ionic liquid aerosol



Hypergolic  
Ionic liquid aerosol ‘reacted’

+  $h\nu$


Reaction dynamics  
and kinetics



C. J. Koh, et al., *Soft ionization of thermally evaporated hypergolic ionic liquid aerosols*, J. Phys. Chem. **115**, 4630 (2011).




# Ionic liquid aerosol generation



- Tunable light source


Hypergolic Ionic liquid aerosol

- size distribution





# Experimental setup









# Identification of reaction products


- photoionization efficiency curve measured








# aerosol kinetics data






# *MD simulations of IL/oxidizer interphases: liquid-liquid*

**t=0**




**t=6ns**




**in liquid/liquid systems, mixing is rapid!**






**For the liquid-liquid interphase intermixing occurs fast (few ns time scale for investigated dimensions).**




# *MD simulations of IL/oxidizer interphases: liquid-gas*



**t=0**



**t=6ns**








**At the liquid-gas interphase HNO<sub>3</sub> quickly adsorbs to the surface and then slowly diffuses to the bulk.**

**There is a certain concentration of HNO<sub>3</sub> at the surface that system would like to maintain to reduce the surface tension**





# Conclusions

- Molecular dynamics can greatly improve our understanding of reactivity of ionic liquids:
  - ID thermal decomposition mechanisms
  - surface tension/diffusion limited processes
- Tunable VUV-PI-TOFMS a powerful experimental tool:
  - Direct detection of ion pairs upon vaporization
  - ID products by mass *and* ionization potential
  - Aerosols make ionic liquids more accessible by MS
- Design a better green rocket fuel with ionic liquids !!!







# Acknowledgements



- Jianbo Liu  
(CUNY Queens College)
- Dmitry Bedrov  
(Wasatch Molecular)
- Schatz Group:
  - Scott Yockel
- Hase Group:
  - Swapnil Kohale
- Leone Group  
(UC Berkeley and ALS):
  - Christine Koh
  - Amir Golan
  - Denisia Popolan
  - Oleg Kostko
  - Chris Harmon
  - Daniel Strasser
- Ionic Liquids Group  
(AFRL Edwards):
  - Gammy Vaghjiani
  - Jerry Boatz
  - Capt. Adam Schenk
  - Lt. Anna Sheppard

**Mike Berman**





# *Questions?!*



